



CME POST-TEST

All post-tests must be completed and submitted online.

EXPIRATION DATE: JANUARY 2024

Earn Category I CME Credit by reading both CME articles in this issue, reviewing the post-test, then taking the online test at <http://cme.aapa.org>. Successful completion is defined as a cumulative score of at least 70% correct. This material has been reviewed and is approved for 1 hour of clinical Category I (Preapproved) CME credit by the AAPA.

LUNG CANCER SCREENING

1. What percentage of patients who present with lung cancer receive a diagnosis of locally advanced metastatic disease?
 - a. 37%
 - b. 59%
 - c. 79%
 - d. 97%
2. What percentage of patients eligible for low-dose CT under the USPSTF guidelines receive screening?
 - a. 2%
 - b. 7%
 - c. 16%
 - d. 23%
3. The 2021 USPSTF recommendations for annual lung cancer screening with low-dose CT include
 - a. all patients with at least a 30 pack-year smoking history.
 - b. patients ages 55 to 80 years with a 30 pack-year smoking history and who are current smokers or have quit in the past 15 years.
 - c. patients ages 55 to 80 years with a 30 pack-year smoking history and who are current smokers or have quit in the past 20 years.
 - d. patients ages 50 to 80 years with a 20 pack-year smoking history and who are current smokers or have quit in the past 15 years.
4. What percentage of patients who participate in annual lung cancer screening with low-dose CT are diagnosed with lung cancer?
 - a. 0.01% to 0.03%
 - b. 0.1% to 0.3%
 - c. 1% to 3%
 - d. 11% to 13%
5. The NLST found what percentage relative reduction in mortality from lung cancer with annual low-dose CT compared with chest radiography?
 - a. 15%
 - b. 20%
 - c. 25%
 - d. 30%

LYME DISEASE COMPLICATIONS

6. Complications in the early disseminated and late phases of Lyme disease include
 - a. Lyme carditis, Lyme arthritis, and Lyme neuroborreliosis.
 - b. Lyme carditis, Lyme arthritis, and Lyme dermatitis.
 - c. Lyme carditis, Lyme arthritis, and Reiter syndrome.
 - d. Lyme carditis, Lyme arthritis, and Lyme conjunctivitis.
7. What is the actual incidence of Lyme disease in the United States?
 - a. 30,000 cases annually
 - b. 47,000 cases annually
 - c. 300,000 cases annually
 - d. 476,000 cases annually
8. What is the most common complication of late Lyme disease?
 - a. Lyme carditis
 - b. Lyme neuroborreliosis
 - c. Lyme arthritis
 - d. uveitis
9. What percentage of patients treated for Lyme disease are diagnosed with PTLDS?
 - a. less than 5%
 - b. up to 20%
 - c. more than 50%
 - d. 62%
10. In regions that are highly endemic for Lyme disease, prophylactic treatment may be warranted using a single dose of
 - a. amoxicillin.
 - b. metronidazole.
 - c. levofloxacin.
 - d. doxycycline.