



DEBUGGING SEPSIS

Andrew Walker, PSM, MS, PA-C
Instructor in Medicine
September 2023

DISCLOSURES

1. I have no relevant financial or commercial interests.
2. Off label use of products denoted by †
3. Our understanding of sepsis and its best treatment is continuing to change rapidly. 2021 SSC updated guidelines in brackets.

OBJECTIVES



Describe the pathophysiology of sepsis



Recognize sepsis early and initiate appropriate therapy quickly




Discuss novel approaches to sepsis treatment

CASE PRESENTATION

- A 68yo M with a hx of HTN, HLD, and alcohol abuse presented to the ED via EMS with abdominal pain.
- He was drowsy, confused, peripherally cold, and cyanotic. His BP was 75/50 with a HR of 125 BPM.
- What next?


E.D. COURSE


- His abdomen was tight and distended and an EKG showed:




- He received 1L of NS and a CT of the abdomen showed extraluminal gas & free fluid consistent with

BEDSIDE ULTRASOUND





During laparotomy, gross fecal peritonitis from a perforated sigmoid colon was confirmed. Resection of the sigmoid colon with closure of the rectal stump and creation of an end colostomy was performed with extensive peritoneal washout. He was transferred to the ICU intubated and still under anesthesia with FiO_2 of 0.4.



BEDSIDE ULTRASOUND

- Blood pressure 88/52 mmHg
- CVP 7 mmHg
- Temperature 35.6°
- ABG pH 7.22 with pCO₂ 25
- WBC 15k
- Lactate 3.0 mmol/L
- Diagnosis?



TREATMENT

Fluids?

Antibiotics?

Sodium
bicarbonate?

Beta-blocker?

Calcium
channel
blocker?

Mechanical
device?

Steroids?

Vitamin C?

Thiamine?

Statin?

Vasopressors?

Inotropes?

NOW WHAT?




He's treated with broad spectrum antibiotics, vasopressors, and is successfully weaned off mechanical ventilation over the course of three days. He's transferred to the floor and suddenly develops shortness of breath.



SaO₂ 82% on pulse oximetry. He's tachycardic and tachypneic. SBP 84/35.

BESIDE ULTRASOUND

- Diagnosis?








EPIDEMIOLOGY

Scope of the burden


- Approximately 31 million people develop sepsis each year globally.
- 6 million deaths/year.
- Increased incidence due to increased chemotherapy, immunosuppression, transplantation, chronic health conditions, and coding.

SEPSIS INCIDENCE

- Today >1.7 million people/yr in the U.S. acquire sepsis and >270,000 will die.
- Claims-based vs EHR-based data
 - Which would you suspect is greater?
 - What are the trends in death?
- Mortality appears to be improving but data are conflicting.
- [Discuss goals of care and prognosis]



PATHO- PHYSIOLOGY



- Pro-inflammatory activation
- Mitochondrial, endothelial, and coagulation dysfunction
- Immunosuppression, oxidative stress, and cellular necrosis leads to MODS and death

An aerial photograph of a city with numerous skyscrapers and buildings. A large green rectangular area is overlaid on the left side of the image, and a white rectangular frame is overlaid on the right side. In the center, a dark grey rectangular box contains the text.

HISTORY

How did we get here?

BACKGROUND

- “Hectic fever, at its inception, is difficult to recognize but easy to treat; left unattended it becomes easy to recognize and difficult to treat.”
 - Niccolò Machiavelli
- Dr. Rivers pioneering work in 2001 EGDT showed a 16% absolute reduction in hospital mortality.
- Aftermath


Early recognition

Early volume
resuscitation

Early
antibiotics

BACKGROUND CONT.

- Previously Sepsis = Infection + 2 or more SIRS criteria
 - HR >90, Temp >38.3, RR >20, WBC >12k
- One in eight patients with severe sepsis will be missed using SIRS criteria
- Ideally, we could develop an early detection method and a definitive diagnostic marker.



DEFINITIONS AND GUIDELINES

- Definitions and guidelines continue to evolve (Sepsis-1 in 1991, Sepsis-2 in 2001, and Sepsis-3 in 2016)
 - CMS and ICD-10 continue to differ from Surviving Sepsis Campaign.
 - There is no gold standard “sepsis test” – *it is a syndrome.*
 - Sepsis is “life-threatening organ dysfunction caused by a dysregulated host response to infection. For clinical operationalization, organ dysfunction can be represented by an increase in the SOFA score of 2 points or more, which is associated with an in-hospital mortality >10%.”

DEFINITIONS AND GUIDELINES CONT.

- Septic shock is “a subset of sepsis in which particularly profound circulatory, cellular, and metabolic abnormalities are associated with a greater risk of mortality than with sepsis alone. Patients with septic shock can be clinically identified by a vasopressor requirement to maintain a mean arterial pressure of 65 mm Hg or greater and serum lactate level greater than 2 mmol/L (>18 mg/dL) in the absence of hypovolemia. This combination is associated with hospital mortality rates greater than 40%.”
- Decoupled sepsis from uncomplicated infections meeting SIRS criteria.
- In April 2018, the SSC provided a 1-hour bundle for treatment when diagnosed.
- In November 2021 we received another set of recommendations.

qSOFA SCORE

Q

SBP < 100

RR > 22

GCS < 15



QUICK SOFA (qSOFA)

- SOFA score is an illness-severity score used to predict mortality of critically ill patients.
- Patients with suspected sepsis can be rapidly identified if they meet *at least 2 of three criteria* of the score.
- Lactate is superior to qSOFA for sepsis prognostication.
- **Take home:**
 - SIRS may over AND under diagnose but still has a role to play
 - SOFA is cumbersome in the ED but great for ICU patients
 - qSOFA is NOT a diagnostic tool and [should not be used as a single screening tool]


DEFINITIONS CONT.

	SEP-1	SEP-3
Sepsis	Infection + ≥ 2 SIRS criteria	≥ 2 qSOFA including hypotension
Severe Sepsis	Sepsis + organ dysfunction*	N/A
Septic Shock	Sepsis + refractory hypotension +/- lactate ≥ 4	Vasopressors and lactate > 2 mmol/L

SEPSIS PHENOTYPES

Pheno-type	Characteristics
alpha	Least vasopressors
Beta	Older, more chronic illness and renal dysfunction
Gamma	More inflammation and pulmonary dysfunction
Delta	More cardiovascular and liver dysfunction. Increased septic shock

- Heterogenous syndrome with an overly broad definition
- Different biomarker elevation and mortality
- Directed treatment may improve mortality



PROBLEMS WITH RECENT GUIDELINES

- Several financial conflicts of interest exist, and several strong recommendations are based on weak evidence (e.g lactate, 1hr).
- Various medical societies were not consulted and have refused to endorse them.
- Disregard clinician judgement with fixed time frames and fluid volumes.
- qSOFA is specific but not sensitive for organ dysfunction (96.1 % vs 29.7) and early risk assessment.
- However, checklists and reminders can be beneficial. Hospitals with higher compliance rates have lower mortality.
- Other risk-stratification scores are available (e.g. MEWS and NEWS) to recognize *critical illness*.

Williams JM, Greenslade JH, McKenzie JV, et al. SIRS, qSOFA and organ dysfunction: insights from a prospective database of emergency department patients with infection. *Chest* 2017;151:586-596.

Giamarellos-Bourboulis EJ, Tsaganos T, Tsangaris I, et al. Validation of the new Sepsis-3 definitions: proposal for improvement in early risk identification. *Clin Microbiol Infect* 2017;23:104-9.

RISK FACTORS FOR SEPSIS

1. Advanced age
2. Immunosuppression/steroid use/malnutrition
3. DM/CA/HIV/liver disease
4. Recent abx/drug-resistance
5. Recent procedures or travel
6. Alcohol/drug use



Initial Signs and Symptoms

- Temp >38.3 or $<36^{\circ}\text{C}$
- HR >90 BPM
- RR $>20/\text{min}$




Signs of end organ hypoperfusion

- Warm diaphoretic skin \rightarrow cool, cyanotic, mottled, and delayed capillary refill in shock
- AMS, restlessness, agitation, or obtundation
- Oliguria or anuria
- Ileus or absent bowel sounds

SIRS AND FEVER DIFFERENTIAL

- SIRS

- Sepsis
- Trauma/Burns/Surgery
- Pancreatitis
- Chemical pneumonitis
- Anaphylaxis
- Post operative
- Pulmonary emboli




LABORATORY FINDINGS

- WBC >12,000 or <4,000
- Glucose >140 mg/dL in the absence of diabetes.
- CRP > 2 S.D. above normal
- Arterial hypoxemia
 - p/f <300
- Cr increase > 0.5 mg/dL
- INR >1.5 or aPTT >60s
- Platelet count <100,000
- Total bilirubin >4 mg/dL
- Lactate >2 mmol/L
- Procalcitonin >2 S.D. above normal
- Adrenal insufficiency or euthyroid sick syndrome

LACTATE

- Produced daily by many organs under normal conditions but \uparrow during inadequate O_2 delivery resulting in tissue hypoxia and anaerobic metabolism.
- Elevated in $\sim 2/3$ of pts with severe sepsis or septic shock & $\infty \uparrow$ mortality rate.
- However, sepsis results in an *impaired ability* of the tissues to extract O_2 and \uparrow delivery does not entirely reverse lactate formation.
- Using cap refill may be just as good (if not better) to guide resuscitation.
- Other causes of hyperlactatemia:
 - Hepatic dysfunction / \downarrow clearance
 - Cancer
 - Hyperadrenergia
 - Thiamine deficiency
 - Toxins/drugs/ethanol
 - Ketoacidosis
 - Inborn errors of metabolism



1. Bandarn Suetrong, MD; and Keith R. Walley, MD. Lactic Acidosis in Sepsis: It's Not All Anaerobic: Implications for Diagnosis and Management. CHEST 2016; 149(1):252-261
2. Casserly B, Phillips GS, Schorr C, et al. Lactate measurements in sepsis-induced tissue hypoperfusion: results from the Surviving Sepsis Campaign Database. Crit Care Med. 2015;43(3):567-573.
3. Hernandez G et. al. Effect of a Resuscitation Strategy Targeting Peripheral Perfusion Status vs Serum Lactate Levels on 28-Day Mortality Among Patients With Septic Shock: The ANDROMEDA-SHOCK Randomized Clinical Trial. JAMA. 2019 Feb 19;321(7):654-664. doi: 10.1001/jama.2019.0071.



TREATMENT

GOALS

1. Stabilize hemodynamically
2. Obtain source control
3. Manage host complications

THE SURVIVING SEPSIS CAMPAIGN BUNDLE: 2018 UPDATE

1. Measure lactate (recheck if $>2\text{mmol/L}$)
2. Obtain blood cultures prior to administering antibiotics
3. Administer broad-spectrum antibiotics
4. Rapidly administer 30mL/kg crystalloid for hypotension or lactate ≥ 4
5. Apply vasopressors if hypotensive during or after fluid resuscitation to maintain $\text{MAP} \geq 65$

ANTIBIOTICS



IV administration **ASAP after recognition and within one hour** for sepsis and septic shock (strong recommendation, moderate quality of evidence).



Recommendation based upon data demonstrating \uparrow mortality for every hour of delay in antibiotic administration for infected pts with organ dysfunction and/or shock. *However, some meta-analyses report no benefit of rapid antibiotic administration.*




It is currently unclear if antibiotic administration within one hour is better than within three in sepsis compared to septic shock.






Current CMS guidance is to administer antibiotics ASAP and within 3 hours of sepsis or septic shock diagnosis.

ANTIBIOTICS

- Start early
 - Each hour delayed \uparrow 1-year mortality by 10%
- Start broad for likely source and base on any prior susceptibility data
- Utilize pharmacists to ensure appropriate dosing
- Practice good stewardship
- Deescalate ASAP [“shorter” duration]



OBTAIN SOURCE CONTROL



INTRAVENOUS FLUID

- Volume
 - Guidelines suggest 30 mL/kg
 - Document reasons for deviation
 - How do we measure response?
 - A positive daily fluid balance is strongly associated with increased mortality
 - CLOVERS trial did not show difference in mortality with restrictive fluid strategy


INTRAVENOUS FLUID CONT.

- Type

- **Physiologically balanced solutions** make physiologic sense but have failed to demonstrate a definitive decrease in mortality
 - SALT-ED trial did not show a ↓ in hospital LOS but ↓ major adverse kidney events within 30 days compared to NS
 - SMART trial showed balanced crystalloids ↓ death and renal dysfunction compared to NS. Non-blinded, single-center study of critically ill
- When choosing colloid, **choose albumin** particularly if 3rd spacing is present
- Cost vs benefit
- More trials pending but [use balanced crystalloids instead of NS]

INOTROPES

- Patients with septic shock may develop impaired myocardial function.
- The pathophysiology of this “septic cardiomyopathy” is not fully established.
- Patients may benefit from inotropic support (e.g., dobutamine) but targeting a specific SvO₂ within a specific timeframe does not improve outcomes.





STEROIDS

- Indicated only for patients with septic shock refractory to fluids and vasopressors
- Stress dosing according to studies is 50mg hydrocortisone q6 hrs or the equivalent
- No need to perform an ACTH
- Tapering not necessary if used for short duration.
 - Consider tapering when vasopressors are no longer needed.


VITAMIN C†




- Rooted in biologic rationale.
 - Key cofactor in endothelial function and catecholamine synthesis
- Headlines vs data
 - 2016 retrospective before-after study of 94 patients (half received placebo) in a single ICU in Virginia.
 - **HAT** = hydrocortisone, ascorbic acid (1.5g IV q6h), and thiamine.
 - Retrospective before-after study comparing mortality over 7 months with those treated showed a decrease from 40.4% to 8.5%!
 - Not controlled, lots of exciting results but follow up research has shown no benefit or even harm (*e.g. J. Clin. Med. 2019, 8(4), 478; or Crit Care Med. 2019 Jun;47(6):774-783, Crit Care Med. 2020 July, 48(7) p e620-e628, and JAMA. 2021;325(8):742-750 VICTAS RCT*)

DIFFERING RESULTS

Marik



Litwak



FECAL MICROBIOTA TRANSPLANT†

- Gut microbiota serves as a physical barrier and immune modulator with disruption leading to extraintestinal disease.
- FMT may be used to reestablish the normal microbial system if dysbiosis and reduced bacterial variability occur due to steroids, sepsis, and/or antibiotics.
- FDA released a warning 2019 about the risk of MDR organisms being transplanted but in 2022 and 2023 approved a rectal suspension then an oral capsule (SER-109;Vowst) for adults with more than one episode of C.diff respectively.
- Utilize *caution* when introducing a high antigenic load in the setting of increased membrane permeability.


Wurm P, Spindelböck W, Krause R, et al. Antibiotic-associated apoptotic enterocolitis 1 in the absence of a defined 2 pathogen: The role of intestinal microbiota depletion.

Crit Care Med. 2017 Jun; 45(6): e600–e606.

Wei et al. Critical Care Successful treatment with fecal microbiota transplantation in patients with multiple organ dysfunction syndrome and diarrhea following severe sepsis (2016) 20:332 DOI 10.1186/s13054-016-1491-2

Kassam Z, et al. Fecal microbiota transplantation for Clostridium difficile infection: systematic review and meta-analysis. Am J Gastroenterol. 2013; 108(4):500.



OTHERS†



- Antibodies
 - anti-endotoxin, anti-enterobacteriaceae, anti-TNF, adrecizumab
- Antagonists
 - IL1, TLR-4, TNF receptor, bradykinin
- Anti-inflammatories/antioxidants
 - N-acetylcysteine, NO inhibitors, ibuprofen, selenium, HAT
- Others
 - G-CSF, antithrombin, tifacogin, GH, calcitriol, levosimendan, hypothermia, hyperoxia, HTS, angiotensin II, alkaline phosphatase, recombinant human soluble thrombomodulin, adrenomedullin, angiotensin II, InnovoSep (cilengitide), targeted antibodies, etc.

NOVEL/EUA APPROACHES[†]

- Cytokine, receptor, micro-RNA, and various proteins as rapid diagnostic biomarkers
- Inflammatory molecule filtration
- Bacteriophages to immune checkpoint inhibitors
- Vaccines
 - Injected or implanted under the skin
 - Combines immunogenic antigens from multiple pathogens and immune cell-recruiting biomaterial scaffolds
- AI



<https://www.seas.harvard.edu/news/2021/07/biomaterial-vaccines-ward-broad-range-bacterial-infections-and-septic-shock>

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7805252/>

<https://www.dovepress.com/mir-486-5p-serves-as-a-diagnostic-biomarker-for-sepsis-and-its-predict-peer-reviewed-fulltext-article-JIR>

TX SUMMARY

Therapy	Specifics	Pearls
Initial resuscitation	30cc/kg in first hour?	Consider LR/albumin
	MAP \geq 65 mmHg	PLR, cap. refill, lactate
Antibiotics	Initiate broad spectrum [including fungal if high risk]	E.g. vanc/pip-tazo Consider procal [to stop]
	Obtain source control	
Steroids	Only if septic shock refractory to fluids/vasopressors	
Vasopressors	NE then VP [then Epi]	Avoid dopamine for most [Peripherally OK if critical]
VTE prophylaxis	[Lovenox instead of heparin]	

TAKE HOME POINTS

1. Sepsis is a life-threatening response to an infection that must be diagnosed early.
2. Sepsis should be treated quickly based on protocols with IV fluids and broad-spectrum antibiotics while incorporating clinical expertise for personalized care.
3. Source control must be obtained.
4. Novel diagnostic markers and therapeutics are needed to improve patient outcomes.



THANK YOU

walker.andrew@mayo.edu