

Detoxing and Dopesick

Sarah Willis Tiesing, MSN, FNPC

TeamHealth Hospital Medicine Lead APC

CJW Hospitals, Richmond, VA

Non-Declaration Statement

I have no relevant relationships with ineligible companies to disclose within the past 24 months.

Objectives

01

The learner will be able to identify the signs and symptoms of acute alcohol and opioid withdrawal.

02


The learner will be able to identify the risks and complications of alcohol and opioid withdrawal.

03

The learner will be able to identify best practice pharmacologic therapies for acute alcohol and opioid withdrawal.

Alcohol Abuse and Dependence

- ◇ Prevalence in U.S.
 - ◇ 29.5 million ages 12 and older with AUD
 - ◇ 10.6% in this age group
 - ◇ 11-32% inpatients
 - ◇ Approximately 50% of those with alcohol use disorder experience alcohol withdrawal
 - ◇ 500,000 episodes of withdrawal severe enough to require pharmacologic treatment each year
 - ◇ Excessive alcohol consumption: 3rd leading preventable cause of death in US



Alcohol Abuse and Dependence

- ◇ Risk Factors
 - ◇ Substance abuse history
 - ◇ History of mental illness
 - ◇ History as victim of abuse
- ◇ Complications
 - ◇ High risk behaviors
 - ◇ Drunk driving
 - ◇ Unsafe sex
 - ◇ Falls/trauma
 - ◇ Liver damage, cirrhosis
 - ◇ GERD/PUD
 - ◇ Esophageal varices
 - ◇ Acute Hemorrhage – increased risk of bleeding
 - ◇ Heart Failure
 - ◇ Insulin resistance
 - ◇ Wernicke's encephalopathy

Alcohol Withdrawal

ARE YOU DRUNK?


YES

NO

- ◇ Predictors of withdrawal:
 - ◇ Consumption of more drinks per occasion
 - ◇ Prolonged, sustained intake
 - ◇ Presence of more alcohol-related problems
 - ◇ No withdrawal symptoms >24 hrs after last drink less likely to develop symptoms

Alcohol Withdrawal

- ◊ Alcohol Intoxication:
CNS depressant
- ◊ Alcohol Withdrawal:
CNS hyperactivity




Minor Alcohol Withdrawal

- ◇ Insomnia
- ◇ Tremulousness
- ◇ Mild anxiety
- ◇ GI upset, anorexia
- ◇ Headache
- ◇ Diaphoresis
- ◇ Palpitations



**IF ALCOHOL CAN DAMAGE
YOUR SHORT TERM MEMORY**

A man with a beard, wearing a dark blue sailor's uniform with white stripes on the collar, is smiling and holding a glass of dark beer. The background is a blurred bar setting.

**IMAGINE THE DAMAGE
ALCOHOL CAN DO**

Alcohol Withdrawal


◇ 20% of patients experience moderate to severe symptoms:

◇ Hallucinosi

◇ Seizures

◇ Delirium tremens

Alcohol withdrawal spectrum



Timing of alcohol withdrawal syndromes

Syndrome	Clinical findings	Onset after last drink
Minor withdrawal	Tremulousness, mild anxiety, headache, diaphoresis, palpitations, anorexia, GI upset; Normal mental status	6 to 36 hours
Seizures	Single or brief flurry of generalized, tonic-clonic seizures, short post-ictal period; Status epilepticus rare	6 to 48 hours
Alcoholic hallucinosis	Visual, auditory, and/or tactile hallucinations with intact orientation and normal vital signs	12 to 48 hours
Delirium tremens	Delirium, agitation, tachycardia, hypertension, fever, diaphoresis	48 to 96 hours

Alcoholic Hallucinosis

- ◇ Typically begins 12-24 hrs after last drink
- ◇ Resolves in another 24-48 hrs
- ◇ Usually visual
- ◇ NO disorientation
- ◇ Vital signs are normal
- ◇ Contributing factors:
 - ◇ Possibly genetic
 - ◇ Possibly due to reduced thiamine absorption

Alcohol Withdrawal Seizures

- ◇ Occur within 6-48 hrs after cessation or after significant reduction
- ◇ Occurs in up to 10-30% of withdrawal patients
- ◇ Typically tonic-clonic, brief and single or in short cluster of 2-3
- ◇ Potential for status epilepticus is low
- ◇ Risk factors:
 - ◇ Concurrent withdrawal from benzos or sedative-hypnotics
 - ◇ Hypokalemia
 - ◇ Thrombocytopenia
 - ◇ Repeat withdrawals – “kindling effect”
 - ◇ Ages 50-60 yrs may emerge

Delirium Tremens

- ◇ Withdrawal Delirium
- ◇ Begins between 72-96 hours after last drink
- ◇ Up to 5 % of hospitalized withdrawal patients
- ◇ Risk factors:
 - ◇ Increasing age
 - ◇ Concurrent illness
 - ◇ w/drawal w/+ BAC
- ◇ Rapid-onset
- ◇ Disorientation
- ◇ Hallucinations
- ◇ Agitation
- ◇ Autonomic hyperactivity
 - ◇ Fevers
 - ◇ Tachycardia
 - ◇ HTN
 - ◇ Diaphoresis

Assessment Tools

- ◇ CIWA-Ar score
- ◇ RASS score
- ◇ Moderate and Severe Alcohol Withdrawal
Rapid Overview (uptodate)

CIWA-Ar Scale

Clinical Institute Withdrawal Assessment Scale for Alcohol, Revised

Clinical Institute Withdrawal Assessment for Alcohol – revised (CIWA-Ar) scale

Clinical Institute Withdrawal Assessment for Alcohol revised	
Symptoms	Range of scores
Nausea or vomiting	0 (no nausea, no vomiting) – 7 (constant nausea and/or vomiting)
Tremor	0 (no tremor) – 7 (severe tremors, even with arms not extended)
Paroxysmal sweats	0 (no sweat visible) – 7 (drenching sweats)
Anxiety	0 (no anxiety, at ease) – 7 (acute panic states)
Agitation	0 (normal activity) – 7 (constantly trashes about)
Tactile disturbances	0 (none) – 7 (continuous hallucinations)
Auditory disturbances	0 (not present) – 7 (continuous hallucinations)
Visual disturbances	0 (not present) – 7 (continuous hallucinations)
Headache	0 (not present) – 7 (extremely severe)
Orientation/clouding of sensorium	0 (orientated, can do serial additions) – 4 (Disorientated for place and/or person)

- ◇ Score:
- ◇ <10: Very mild withdrawal
- ◇ 10 to 15: Mild withdrawal
- ◇ 16 to 20: Modest withdrawal
- ◇ >20: Severe withdrawal



**DAD JOKES ABOUT
ALCOHOL WITHDRAWAL ARE SHAKY**

CIWA I DID THERE?

Richmond Agitation Sedation Scale (RASS)

Richmond Agitation-Sedation Scale

	Target RASS Value	RASS Description
+4	Combative	Combative, violent, immediate danger to staff
+3	Very Agitated	Pulls or removes tube(s) or catheter(s); aggressive
+2	Agitated	Frequent non-purposeful movement, fights ventilator
+1	Restless	Anxious, apprehensive but movements are not aggressive or vigorous
0	Alert and Calm	
-1	Drowsy	Not fully alert, but has sustained awakening to voice (eye opening & contact greater than 10 seconds)
-2	Light Sedation	Briefly awakens to voice (eye opening & contact less than 10 seconds)
-3	Moderate Sedation	Movements or eye opening to voice (but NO eye contact)
-4	Deep Sedation	No response to voice, <u>but</u> has movement or eye opening to physical stimulation
-5	Unarousable	No response to voice or physical stimulation

Alcohol withdrawal treatment

Alleviate symptoms, prevent complications

Identify and correct underlying metabolic derangements

- Benzodiazepines
- IV fluids
- MVI/Thiamine/Folate
- Monitor with CIWA-AR and Vital signs

Alcohol Withdrawal Pharmacologic Therapy

- ◇ Benzodiazepines are first-line treatment
 - ◇ Treat agitation
 - ◇ Prevent progression of withdrawal
 - ◇ Diazepam (Valium), Lorazepam (Ativan), Chlordiazepoxide (Librium)
 - ◇ Diazepam and Chlordiazepoxide generally preferred
 - ◇ Lorazepam preferred for advanced cirrhosis or acute alcoholic hepatitis



Symptom-Triggered Benzo Dosing

- ◇ Valium dosing
 - ◇ 5-20mg PO/IV as needed using CIWA scale – every 1-4 hrs
- ◇ Lorazepam dosing
 - ◇ 2-4mg PO/IV as needed using CIWA scale – every 1-4 hrs

Phenobarbital Dosing

- ◇ Off-label use
- ◇ Acute withdrawal:
 - ◇ Initial dose 130mg or 260mg x 1
 - ◇ Then 130mg every 15-30 min as needed
- ◇ Maintenance dosing
 - ◇ 130-260mg/day in 2-3 divided doses for 3-5 days, with taper of 10% reduction/day.

Clonidine

- ◇ Off-label use
- ◇ Similar mechanism of action to Precedex
- ◇ Stop all home BP meds
- ◇ Max Clonidine out as much as BP will allow
- ◇ Will treat sympathetic nervous system response
 - ◇ Decrease BP/HR
 - ◇ Decrease agitation

Front-loading vs. Symptom-based treatment

◇ Front-loading

- ◇ Is patient likely to be harmed by prolonged periods of hypertension and tachycardia or by complications from prolonged physical restraint?
- ◇ Rapidly performed
- ◇ Heavier sedation
- ◇ Needs ICU admission
- ◇ Diazepam 5-10mg Q10-15min
 - ◇ Up to 500mg
- ◇ Lorazepam 2-4mg IV Q15-20min
- ◇ Can also use Chlordiazepoxide 25-100mg Q1H

Symptom-based treatment

◇ Symptom-Triggered

- ◇ Use CIWA-Ar every 10-15 min-6 hrs
- ◇ RASS for intubated patients or for manipulative patients
- ◇ Requires less medication and shorter treatment period than fixed schedule
- ◇ Usually benzos

Refractory DT treatment

Resistant alcohol withdrawal

ICU admission

Phenobarbital with benzodiazepine – work synergistically

- Generally limit to 15mg/kg in first 24 hrs
- Maintenance dose 130-260mg/day divided for 3-5 days, 10% reduction/day thereafter
- Not enough research to recommend as a single agent – but could be recommended in future

Precedex and Propofol – need further research

Dopesick



Opioid Use Disorder (OUD)

- ◆ *****Chronic, relapsing illness*****
- ◆ Epidemic proportions
- ◆ In 2021, 9.2 million people age 12 or older misused opioids in the U.S.
- ◆ Between 2002 and 2018, heroin use and disorder nearly doubled in the U.S.
- ◆ 106,699 drug overdose deaths in the U.S. in 2021
 - ◆ 75% involved an opioid- 80,411 deaths
 - ◆ >130 people die from opioid-related drug overdoses per day
 - ◆ Increasing due to cutting w/Fentanyl

Opioid Use Disorder

◇ Risk factors


- ◇ Hx substance use disorder
- ◇ Younger age
- ◇ More severe pain
- ◇ Mental disorders
- ◇ Hx childhood maltreatment

◇ Complications

- ◇ Infection – cellulitis, IE, OM
- ◇ HIV, Hepatitis, TB, syphilis
- ◇ Narcotic bowel syndrome
- ◇ Hyperalgesia
- ◇ Overdose

Opioids and the Human Brain

- ◇ Opioids interfere with the way that the neurons either send, receive or process their neurotransmitters.
 - ◇ Heroin activates the neurons
 - ◇ Heroin acts as a “fake” neurotransmitter
- ◇ Natural vs. Drug rewards
 - ◇ Drug use = fewer neurotransmitters in reward circuit
 - ◇ Fewer receptors to receive signals
 - ◇ Contributes to tolerance



“Skin-Popping”



Opioid Withdrawal

- ◇ Craving
- ◇ Dysphoria, restlessness
- ◇ Watering eyes, runny nose
- ◇ Body aches, joint pain
- ◇ Nausea, vomiting, abdominal cramping, diarrhea

COWS Scale	
Pulse	GI upset
Sweating	Tremor
Restlessness	Yawning
Pupil size	Anxiety Irritability
Bone/joint aches	Goosebumps
Runny nose/tearing	

Acute Opioid Withdrawal Treatment

Buprenorphine and Methadone for natural-occurring withdrawal only - do not use in iatrogenic withdrawal

Fluid resuscitation

Adjunctive medications

- Anti-emetics - Zofran, Compazine, Phenergan
- Anti-diarrheal - Loperamide, Octreotide
- Anti-anxiety, muscle relaxant - Diazepam
- Alpha-2 adrenergic agonists - Clonidine

Clonidine

- ◇ Off-label use Opioid withdrawal
 - ◇ Adjunct to opioid agonist for relief of withdrawal symptoms.
 - ◇ Can be used as primary treatment
- ◇ Dosing
 - ◇ Again, stop all BP meds and use max dose
 - ◇ Initial dose 0.1-0.2mg depending on BP/HR
 - ◇ Then 0.1-0.3mg every 6-8hrs based on COWS score or symptom severity

Medication- assisted treatment (MAT)

“Recovery status is best defined by factors other than medication status. Neither medication-assisted treatment of opioid addiction nor the cessation of such treatment by itself constitutes recovery. Recovery status instead hinges on broader achievements in health and social functioning – with or without medication support.”

- Recovery experts A. Thomas McLellan and William White

MAT

- ◇ Reduce morbidity and mortality
- ◇ Decrease OD deaths
- ◇ Reduce transmission of infectious disease
- ◇ Increase treatment retention
- ◇ Improve social functioning
- ◇ Reduce criminal activity

MAT

Buprenorphine and Methadone

- ◇ Acute pain Indications
 - ◇ Can continue maintenance med (if on Buprenorphine only) although not generally effective in practice
 - ◇ Maximize non-pharmacologic therapies
 - ◇ Supplement with additional opioid
 - ◇ Always d/c Naltrexone
- ◇ Use for naturally-occurring withdrawal
 - ◇ Methadone 10mg IM or 20mg PO
 - ◇ Better for patients addicted to high doses opioids
 - ◇ Buprenorphine 4-8mg SL
 - ◇ Risk of dental decay noted in 2022

Buprenorphine vs. Methadone

Buprenorphine

- ◇ Generally preferred for treatment over Methadone
- ◇ Initiate in a state of mild to moderate withdrawal
- ◇ Ceiling effect
- ◇ Combined with Naloxone to help prevent IV abuse
 - ◇ Naloxone has poor SL bioavailability
 - ◇ Suboxone, Zubsolv branded names

Methadone

- ◇ Full mu-opioid agonist
- ◇ Not safe in overdose
 - ◇ Additive effect with other opioids

Buprenorphine Dosing

- ◇ Start when signs opioid w/drawal occur, but no sooner than 12 hours after last opioid use.
 - ◇ ER Injection - SubQ, 100-300mg once monthly
 - ◇ SL tablet/film
 - ◇ 4 - 8 mg initial dose (8-32mg seen in ED)
 - ◇ Maintenance: adjust in increments of 4mg to a level that maintains treatment and suppresses opioid w/drawal treatment
 - ◇ Doses >16mg not much better efficacy
 - ◇ Max dose 24mg/day in VA
 - ◇ Dosing frequency - patient preference

Methadone Dosing

- ◇ Overall more risky drug
 - ◇ High abuse liability, resp. depression, QT prolongation, additive effects with benzo and other CNS depressants
- ◇ For acute withdrawal, short-term medically supervised
- ◇ Titrate to approx. 40mg/day in divided doses for 2-3 days
 - ◇ After 2-3 days, gradually decrease dose on a daily basis or at 2-day intervals

MAT SAMHSA Guidelines

<https://store.samhsa.gov/product/Advisory-Sublingual-and-Transmucosal-Buprenorphine-for-Opioid-Use-Disorder-/SMA16-4938>

Xylazine

“Tranq” “Zombie Drug”

- ◇ Alpha-2 Agonist Vet medication – animal sedation and analgesia
- ◇ Commonly laced in heroin, fentanyl, cocaine
- ◇ OD: coma, apnea, bradycardia, hypotension
- ◇ Severe necrotic skin ulcerations and abscesses with parenteral use
- ◇ Naloxone ineffective as reversal agent
- ◇ Treatment: Supportive care



Clinical Pearls

- ◇ Alcohol withdrawal is best managed by symptom-triggered benzodiazepine therapy.
- ◇ Escalating alcohol withdrawal should include combination Phenobarbital and benzodiazepine therapy.
- ◇ Opioid abuse is a chronic relapsing illness.
- ◇ Chronic treatment of opioid use disorder best practice is Buprenorphine and Methadone.
- ◇ Xylazine is an emerging health threat in the US.

References

- Buprenorphine and Naloxone (n.d.) Buprenorphine and Naloxone: Drug information. Retrieved March 10, 2023 from Uptodate database: <https://www.uptodate.com/contents/buprenorphine-and-naloxone-drug-information>
- Centers for Disease Control and Prevention (2022, July 29). *Death rates maps and graphs*. <https://www.cdc.gov/drugoverdose/deaths/index.html>
- Clonidine (n.d.) Clonidine: Drug Information. Retrieved March 10, 2023 from Uptodate database: <https://www.uptodate.com/contents/clonidine-drug-information>
- Curious Tendency. (2011). Neurotransmitter graphic table. Retrieved on June 10, 2022 from - <https://www.pinterest.com/pin/202591683212533294/>
- GPNotebook. (2020). *Clinical Institute Withdrawal Assessment for Alcohol (CIWA-A) scale*. [Infographic]. GPNotebook.com. <https://gpnotebook.com/simplepage.cfm?ID=x2020100717362255460>
- Health Resources and Services Administration (2022). *Opioid crisis*. Retrieved on July 29, 2022 from <https://www.hrsa.gov/opioids>
- Hoffman, R. & Weinhouse, G. (2021). Management of moderate and severe alcohol withdrawal syndromes. In J. Grayzal (Ed.), *Uptodate*. Retrieved on June 8, 2022 from https://www.uptodate.com/contents/management-of-moderate-and-severe-alcohol-withdrawal-syndromes?search=alcohol%20withdrawal&source=search_result&selectedTitle=1~150&usage_type=default&display_rank=1
- Knowyourmeme. (2022). Are you drunk meme. [Image]. Knowyourmeme.com. [Are you drunk? | Check Yes or No | Know Your Meme](https://www.knowyourmeme.com/meme/are-you-drunk/)

References

- Lafayette, R. Azbilliards Forums. (2008). If alcohol can damage your short term memory imagine the damage alcohol can do. [Image]. Forums.azbilliards.com. [Funny pic/gif thread... | Page 2095 | AzBilliards Forums](#)
- Methadone (n.d.) Methadone: Drug information. Retrieved March 10, 2023 from UpToDate database: <https://www.uptodate.com/contents/methadone-drug-information>
- Nasky, K. [@psypharmacopeia] (2018, May 19). Ativan meme. [Tweet]. Twitter. <https://twitter.com/psypharmacopeia/status/997938938882002944>
- ◇ National Institute on Drug Abuse: Advancing Addiction Science (2020). *Drugs, brains and behavior: the science of addiction*. Drugs, Brains, and Behavior: The Science of Addiction: Drugs and the Brain | NIDA (nih.gov)
- ◇ Pace, C. (2022). Alcohol withdrawal: Epidemiology, clinical manifestations, course, assessment, and diagnosis. In M. Friedman (Ed.), *Uptodate*. Retrieved on June 8, 2022, from https://www.uptodate.com/contents/alcohol-withdrawal-epidemiology-clinical-manifestations-course-assessment-and-diagnosis?search=alcohol%20withdrawal&source=search_result&selectedTitle=2~150&usage_type=default&display_rank=2
- Reid, M. (2018, August 11). Review: dopesick by Beth Macy – how to kill 300,000 Americans. *The Times*. <https://www.thetimes.co.uk/article/review-dopesick-by-beth-macy-how-to-kill-300-000-americans-kqj2nwkdh>
- Sevarino, K. (2022). Medically supervised opioid withdrawal drug treatment for addiction. In M. Friedman (Ed.), *Uptodate*. Retrieved on June 8, 2022 from https://www.uptodate.com/contents/medically-supervised-opioid-withdrawal-during-treatment-for-addiction?search=opioid%20withdrawal%20treatment&source=search_result&selectedTitle=1~150&usage_type=default&display_rank=1

References

- ◇ Sevarino, K. (2022) Opioid withdrawal in adults: clinical manifestations, course, assessment and diagnosis. In M. Friedman (Ed.), *Uptodate*. Retrieved on June 8, 2022 from https://www.uptodate.com/contents/opioid-withdrawal-in-adults-clinical-manifestations-course-assessment-and-diagnosis?search=opioid%20withdrawal%20treatment&source=search_result&selectedTitle=3~150&usage_type=default&display_rank=3
- Stolbach, A. & Hoffman, R. (2021). Opioid withdrawal in the emergency setting. In M. Ganetsky (Ed.), *Uptodate*. Retrieved on June 8, 2022 from https://www.uptodate.com/contents/opioid-withdrawal-in-the-emergency-setting?search=opioid%20withdrawal%20treatment&source=search_result&selectedTitle=2~150&usage_type=default&display_rank=2
- Worsham, A. (2014, August 7). *Alcohol Withdrawal*. [Powerpoint slides]. Slideserve. <https://www.slideserve.com/ailsa/alcohol-withdrawal-5446323>
- Zazzle. ImageQ. (2022). *Prohibition we want beer protest c. 1932 poster*. Retrieved on June 9, 2022 from [https://www.zazzle.com/prohibition we want beer protest c 1932 poster-228004255087438156](https://www.zazzle.com/prohibition_we_want_beer_protest_c_1932_poster-228004255087438156)

Questions?

Feel free to contact me at
Sarah_Tiesing@teamhealth.com